

## The Influence of Temperature on Gravitation.

A. K. T. ASSIS<sup>(1)</sup>,(\*) and R. A. CLEMENTE<sup>(2)</sup>

<sup>(1)</sup> *Department of Cosmic Rays and Chronology, Institute of Physics  
State University of Campinas - C.P. 6165, 13081-970 Campinas, SP, Brazil*

<sup>(2)</sup> *Department of Quantum Electronics, Institute of Physics,  
State University of Campinas - C.P. 6165, 13081-970 Campinas, SP, Brazil*

(ricevuto il 4 Gennaio 1993; approvato il 16 Giugno 1993)

**Summary.** — We present some general reasonings using the mass-energy equivalence and Weber's force law for gravitation to show that the weight of a body should increase with its temperature. The two approaches predict a fractional change of weight or, equivalently, of the gravitational constant  $G$ , of one part in  $10^{14}$  per degree.

PACS 12.25 — Models for gravitational interactions.

PACS 01.55 — General physics.

PACS 04.50 — Unified field theories and other theories of gravitation.

According to Einstein's relation,  $E = Mc^2$ , if we increase the energy content of a body, automatically its mass is increased. Using the proportionality between inertial and gravitational masses, we conclude that not only its inertial mass but also its weight should increase if its temperature is raised.

We now calculate this value. To this end, we suppose two spherical solid bodies 1 and 2, with isotropic distributions of mass  $\rho_1(r_1)$  and  $\rho_2(r_2)$ , separated by a distance  $R$ . We will represent their «cold masses», that is, their masses at zero degrees kelvin, by  $M_{10}$  and  $M_{20}$ . From Newton's universal law of gravitation we have at this temperature  $F = -GM_{10}M_{20}/R^2$ . To arrive at temperatures  $T_1$  and  $T_2$  they need to receive a thermal energy. We utilize the law of Dulong-Petit [1] to estimate roughly this amount of energy increase. If body 1 has  $N_1$  and body 2 has  $N_2$  molecules, we obtain  $\delta E_1 = 3N_1 k_B T_1$  and  $\delta E_2 = 3N_2 k_B T_2$ , where  $k_B$  is Boltzmann's constant. Using Einstein's relation we conclude that at these temperatures Newton's law will read

$$(1) \quad F = - \frac{G}{R^2} M_{10} \left( 1 + 3 \frac{N_1 k_B T_1}{M_{10} c^2} \right) M_{20} \left( 1 + 3 \frac{N_2 k_B T_2}{M_{20} c^2} \right).$$

(\*) Also Collaborating Professor at the Department of Applied Mathematics, IMECC, UNICAMP, 13081-970 Campinas, SP, Brazil.

This means that, in general, if we increase the temperatures of bodies 1 and 2 by  $\delta T_1$  and  $\delta T_2$  there will be a fractional change of the gravitational force given by

$$(2) \quad \frac{\delta F}{F} = 3 \left( \frac{k_B \delta T_1}{m_{10} c^2} + \frac{k_B \delta T_2}{m_{20} c^2} \right).$$

In this expression we utilized that  $M_{i0} = N_i m_{i0}$ , where  $m_{i0}$  is the average mass of the molecules constituting body  $i$ .

We now analyse the same problem using Weber's law [2-7]. This is essentially Coulomb's law (or Newton's law) plus terms which depend on the velocity and acceleration between the interacting bodies. There has been a renewed interest in this force law because from it we can derive not only Faraday's law of induction but also Ampère's force between current elements. In the last few years a large number of experiments have been performed trying to distinguish Ampère's force from Grassmann's force [8-11] but this is still an open question [12-13]. When applied to gravitation, Weber's law reads [14]

$$(3) \quad \mathbf{F} = -G m_{10} m_{20} \frac{\hat{\mathbf{r}}_{12}}{r^2} \left[ 1 - \frac{\xi}{c^2} \left( \frac{r^2}{2} - r\dot{r} \right) \right] = \\ = -G m_{10} m_{20} \frac{\hat{\mathbf{r}}_{12}}{r^2} \left[ 1 + \frac{\xi}{c^2} \left( \mathbf{v}_{12} \cdot \mathbf{v}_{12} - \frac{3}{2} (\hat{\mathbf{r}}_{12} \cdot \mathbf{v}_{12})^2 + \mathbf{r}_{12} \cdot \mathbf{a}_{12} \right) \right].$$

In this expression  $\mathbf{F}$  is the force exerted by  $m_{20}$  on  $m_{10}$ ,  $\mathbf{r}_{12} \equiv \mathbf{r}_1 - \mathbf{r}_2$ ,  $r \equiv |\mathbf{r}_{12}|$ ,  $\hat{\mathbf{r}}_{12} \equiv \mathbf{r}_{12}/r$ ,  $\mathbf{v}_{12} \equiv d\mathbf{r}_{12}/dt$ ,  $\mathbf{a}_{12} \equiv d\mathbf{v}_{12}/dt$ ,  $c$  is the light velocity in vacuum and  $\xi = 6$ .

In order to calculate the gravitational interaction between the two previous spheres at temperatures  $T_1$  and  $T_2$ , we make the following: In the first place we suppose each molecule vibrating harmonically around an equilibrium position. For instance, if the equilibrium position of molecule 2 is the origin and that of molecule 1 is  $y_{10}\hat{\mathbf{y}}$ , a possible motion is  $\mathbf{r}_2 = A_2 \sin(\omega_2 t + \theta_2)\hat{\mathbf{x}}$  and  $\mathbf{r}_1 = [y_{10} + A_1 \sin(\omega_1 t + \theta_1)]\hat{\mathbf{y}}$ . For simplicity we suppose  $\omega_1 = n\omega_2$ , where  $n$  is an integer, and  $y_{10}^2 \gg A_1^2 + A_2^2$ , which means that the distance  $R$  between the two spheres is much greater than the amplitudes of oscillation. We then utilize these values of  $\mathbf{r}_1$  and  $\mathbf{r}_2$  in eq. (3) and perform two averages. The first one is on time, integrating from  $t = 0$  to  $t = T_2 = 2\pi/\omega_2$ , and dividing by  $T_2$ , and the second one is on the phases, allowing any value between 0 and  $2\pi$  for  $\theta_1$  and for  $\theta_2$ . We suppose also that  $y_{10}^2 \gg c^2/\omega_1^2 + c^2/\omega_2^2$ . The average value is found to be

$$(4) \quad \langle \mathbf{F} \rangle = -G m_{10} m_{20} \frac{y_{10}\hat{\mathbf{y}}}{|y_{10}|^3} \left( 1 + \frac{\xi}{4} \frac{A_1^2 \omega_1^2}{c^2} \right).$$

We perform this average for the nine combinations of oscillation along the main directions ( $m_1$  oscillating along  $X, Y$  and  $Z$ , and the same for  $m_2$ ). The final average value is found to be

$$(5) \quad \langle \langle \mathbf{F} \rangle \rangle = -G m_{10} m_{20} \frac{y_{10}\hat{\mathbf{y}}}{|y_{10}|^3} \left( 1 + \frac{\xi}{12} \frac{A_1^2 \omega_1^2 + A_2^2 \omega_2^2}{c^2} \right).$$

A molecule in a solid has six degrees of freedom so that  $\langle E \rangle = 6(k_B T/2)$ . As  $\langle E \rangle = 6\langle mv_x^2/2 \rangle$ , we get  $\langle mv_x^2 \rangle = k_B T$  or  $\langle m\mathbf{v} \cdot \mathbf{v} \rangle = 3k_B T$ . In the case under consideration  $\langle v_x^2 \rangle = A^2 \omega^2/2$  so that, applying these results in (5) yields, after integrating for the solid bodies,

$$(6) \quad \langle\langle F \rangle\rangle = -G \frac{M_{10} M_{20}}{R^2} \left[ 1 + \frac{\xi}{6} \left( \frac{N_1 k_B T_1}{M_{10} c^2} + \frac{N_2 k_B T_2}{M_{20} c^2} \right) \right].$$

The fractional change of force according to Weber's law is then given by

$$(7) \quad \frac{\delta \langle F \rangle}{\langle F \rangle} \equiv \frac{\xi}{6} \left( \frac{k_B \delta T_1}{m_{10} c^2} + \frac{k_B \delta T_2}{m_{20} c^2} \right).$$

This result is analogous to (2) and has the same order of magnitude.

To have an idea of this fractional change of gravitational force, which can be considered as a change of weight, of mass or of  $G$ , we take into consideration two solid spheres of iron. Then  $\delta \langle F \rangle / \langle F \rangle = 10^{-14} / ^\circ\text{C}$ . If we change the temperature of the bodies by  $100^\circ\text{C}$ , then  $\delta \langle F \rangle / \langle F \rangle = 10^{-12}$ , which is near to the limit of precision of the modern analytic balances. We propose this experiment in order to know if these effects occur simultaneously or if only the mass-energy equivalence is compatible with the experiments.

In conclusion, we could say that two different ideas and concepts lead to the same conclusion that the gravitational force between two bodies should be a function of their temperatures. Moreover, the expected fractional change of force is of the order  $k_B \delta T / mc^2$ .

Recently, Massa studied some theoretical consequences of a gravitational constant which depends on temperature [15]. In this work we restrict ourselves to pointing out two different models indicating the possible existence of this effect. Other ramifications of this idea are outside the scope of the present work.

It is worthwhile pointing out that Poynting and Phillips examined experimentally a possible dependence of gravitational force on temperature before the coming of Einstein's relation  $E = Mc^2$ , see reference [16]. They showed that if the weight of a body changes due to a modification of its temperature, this change corresponds to less than one part in  $10^{10}$  per degree. There is no contradiction between this result and the effect calculated in this work because we predict a change of only one part in  $10^{14}$  per degree.

\* \* \*

One of the authors, AKTA, wishes to thank Dr. R. de A. Martins for many profitable and stimulating discussions. He thanks also FAPESP and CNPq (Brazil) for financial support during the last years. The authors thank the referee for his valuable suggestion regarding the first version of this work.

## REFERENCES

- [1] F. REIF: *Fundamentals of Statistical and Thermal Physics* (McGraw-Hill, Tokio, 1965), Chapt. 7.

- [2] J. C. MAXWELL: *A Treatise on Electricity and Magnetism*, Vol. 2 (Dover, New York, N.Y., 1954), Chapt. 23.
- [3] A. O'RAHILLY: *Electromagnetic Theory*, Vol. 2 (Dover, New York, N.Y., 1965), Chapt. 11.
- [4] J. P. WESLEY: *Found. Phys. Lett.*, **3**, 443, 471, 586 (1990).
- [5] T. E. PHIPPS jr.: *Apeiron*, **8**, 8 (1990).
- [6] A. K. T. ASSIS and J. J. CALUZI: *Phys. Lett. A*, **160**, 25 (1991).
- [7] A. K. T. ASSIS: *Can. J. Phys.*, **70**, 330 (1992).
- [8] P. GRANEAU: *Nature*, **295**, 311 (1982).
- [9] P. GRANEAU: *Ampère-Neumann Electrodynamics of Metals* (Hadronic Press, Nonantum, 1985).
- [10] P. T. PAPPAS: *Nuovo Cimento B*, **76**, 189 (1983).
- [11] T. E. PHIPPS and T. E. PHIPPS jr.: *Phys. Lett. A*, **146**, 6 (1990).
- [12] P. CORNILLE: *J. Phys. A*, **22**, 4075 (1989).
- [13] P. G. MOYSSIDES: *IEEE Trans. Magn.*, **25**, 4298, 4307, 4313 (1989).
- [14] A. K. T. ASSIS: *Found. Phys. Lett.*, **2**, 301 (1989).
- [15] C. MASSA: *Helv. Phys. Act.*, **62**, 424 (1989).
- [16] J. H. POYNTING and P. PHILLIPS: *Proc. R. Soc. London, Ser. A*, **76**, 445 (1905).